arXiv:1210.5655v1 [hep-th] 20 Oct 2012

Towards an exact density of states in a free conformal field theory

Connor Behan

Department of Physics and Astronomy,

University of British Columbia, Vancouver, BC, V6T 1W9, Canada

(Dated: October 23, 2012)

Abstract

A large class of conformal field theories can be studied by setting the mass to zero in a set of non-interacting fields that form a representation of the Lorentz group. We consider all such field theories defined on a general compact manifold. In the large volume limit, the density of states as a function of energy is shown to obey the same differential equation regardless of the manifold's geometry. From this equation, we compute the leading and next-to-leading behaviour for the density of states, and establish a method that can be iterated to obtain the asymptotic series to any desired precision.

I. INTRODUCTION

The density of states is an example of a thermodynamic quantity which, even in a free field theory, is only ever calculated approximately. Part of the problem is that the notion of a measure $\rho(E)\mathrm{d}E$ giving the number of states with energy in the interval $[E, E + \mathrm{d}E)$ is itself an approximation. For a theory with a discrete spectrum, this assumption is only valid when the energy is high. For this reason, it is natural to approximate the density of states, *asymptotically*. The goal of this work is to better understand the asymptotics of $\rho(E)$ in a free conformal field theory.

Conformal field theories in $1 + 1$ dimensions have been studied extensively since the discovery of worldsheet string theory. More recently, Maldacena's discovery of the AdS/CFT correspondence[1] has ignited an interest in higher dimensional CFTs[2–6] as well. Our results hold for a general d where d is the number of spatial dimensions. For $d \geq 2$, the results appear to be new.

Much is known about the density of states in a two-dimensional CFT ($d = 1$ in our notation). By convention, the volume in such a theory on a circle is usually normalized to 2π. A famous result is the Cardy formula[7] which expresses the density of states in terms of the central charge c (defined so that the vacuum has an energy of $-\frac{c}{12}$). It follows from Cardy's 1986 paper that [1]

$$\rho(E) \propto e^{2\pi\sqrt{\frac{cE}{3}}} = e^{\sqrt{\frac{2c}{3}\pi V E}}$$

for large E. Clearly, this cannot be an exact formula. It does not have an inverse energy as a prefactor for instance. In fact, in 2011, Loran, Sheikh-Jabbari and Vincon showed that

---

[1] For an explanation of the notation, we will use $\propto$ in equations that are roughly true to illustrate a conceptual point. For example $f(x) \propto g(x)$ means that the leading term of f is proportional to the leading term of g. A symbol we use more rigorously is $\sim$. We say that $f(x) \sim g(x)$ if $f(x) = g(x) + o(g(x))$. In other words, the leading terms are equal and as we let $x \to \infty$, $f - g$ becomes negligible compared to f and g.

$$\begin{aligned}\rho(E) &\sim \frac{2\pi^2 c}{3}\frac{I_1\left(2\pi\sqrt{cE/3}\right)}{2\pi\sqrt{cE/3}} \\ &= \frac{\pi c V}{3}\frac{I_1\left(\sqrt{2\pi c V E/3}\right)}{\sqrt{2\pi c V E/3}}\end{aligned}$$

where I_1 is the modified Bessel function of the first kind[8]. Bessel functions have a well known asymptotic series[9] and we can use this series to see that the above formula is consistent with and extends the Cardy formula. For example, the crudest approximation beyond the Cardy formula is

$$\rho(E) \sim \frac{cV}{6(2\pi)^{\frac{1}{4}}(cVE/3)^{\frac{3}{4}}} e^{\sqrt{\frac{2c}{3}\pi V E}} \tag{1}$$

The main property assumed in these derivations is that the partition function is invariant under modular transformations. For our starting point, we instead assume that the partition function is that of a free theory. This allows us to treat the case of higher dimensions in which a manifold is not necessarily flat.

Using dimensional analysis and conformal invariance, there is a simple argument that reveals how $\rho(E)$ must roughly scale with E for a translation invariant system. Energy is extensive so it must be proportional to the volume. For a CFT, the only other scale is the one set by the temperature so by dimensional analysis,

$$E = CVT^{d+1}$$

where C is some dimensionless constant. The same arguments can be applied to entropy, giving us:

$$S = \tilde{C}VT^{d} \propto V^{\frac{1}{d+1}} E^{\frac{d}{d+1}}$$

Finally, Boltzmann's formula relates the entropy to the density of states:

$$\rho(E) \propto e^{S} \propto e^{E^{\frac{d}{d+1}}}$$

The goal of this work is to improve on this formula. If $\zeta(\sigma) = \sum_{n=1}^{\infty} \frac{1}{n^{\sigma}}$ is the Riemann zeta function, $\zeta^{*}(\sigma) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{\sigma}}$ is the alternating zeta function and ω_d is the volume of a unit ball in $\mathbb{R}^d$, the result that will be derived is:

$$\begin{aligned} \rho(E) &\sim C_d a^{\frac{1}{2(d+1)}} E^{\frac{-d-2}{2(d+1)}} e^{\left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{1}{d+1}} a^{\frac{1}{d+1}} E^{\frac{d}{d+1}}} \\ a &= \frac{d!\omega_d[s\zeta(d+1)+s^*\zeta^*(d+1)]V}{(2\pi)^d} \end{aligned} \tag{2}$$

in a theory that has s species of bosons and s^* species of fermions. C_d is a constant depending only on the dimension which can be computed without much difficulty. When only the leading behaviour of the density of states is desired, it is common to see the entropy written instead. Taking the logarithm of our main result,

$$\log \rho(E) \sim \left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{1}{d+1}} a^{\frac{1}{d+1}} E^{\frac{d}{d+1}} - \frac{1}{2(d+1)} \log\left(C_d a E^{d+2}\right) \tag{3}$$

The prefactor in (2) or equivalently the log term in (3) is new and will be referred to as “the logarithmic correction” in this work. The assumptions we make about the physics are captured in the value of a. Once a is known, the form of (2) is generally true. Consider, for example, the maximally supersymmetric large N Yang-Mills theory. This theory is widely studied for its direct role in the AdS/CFT correspondence. The Super Yang-Mills entropy for a coupling parameter of zero

$$S = \left(\frac{256}{27}\frac{\pi^2 N^2 V}{6}\right)^{\frac{1}{4}} E^{\frac{3}{4}}$$

appears frequently in the literature[10–13]. Substituting $d=3$ and $s=s^*=8N^2$ into our formula for a, we see that the entropy above is precisely the first term in (3). We have therefore computed a logarithmic correction to the well known result. In the interacting version, [10–12] have found that the physics can still be well described if the non-interacting a value of $\frac{\pi^2 N^2 V}{6}$ is multiplied by a function of the t'Hooft coupling parameter (which is $\frac{3}{4}$ in the infinite volume limit). Our analysis does not have anything to say about what this funciton might be, but once it is specified, terms in the asymptotic series for $\rho(E)$ can be found using the same method that we apply to a free theory. Another non-free theory in which our result might hold is quantum chromodynamics in the high energy limit. This non-conformal field theory has a scale $\Lambda_{\text{QCD}} \approx 217\text{MeV}$ which emerges at the quantum level. A famous result[14, 15] is asymptotic freedom which states that the running coupling of QCD vanishes at the highest energy scales. Specifically,

$$\alpha(q^2) = \frac{1}{1+\frac{33-2n_{\text{f}}}{12\pi}\log\left(\frac{q^2}{\Lambda^2_{\text{QCD}}}\right)}$$

where q^2 is the squared momentum and n_{f} is the number of fermions. A study of the thermodynamics in this limit[16] has found that the free energy is given by

$$F = -\frac{8\pi^2 V}{45} T^4 \left[\left(1 + \frac{21}{32} n_{\mathrm{f}}\right) - \frac{15}{4}\left(1 + \frac{5}{12} n_{\mathrm{f}}\right) \frac{\alpha(q^2)}{\pi} + O\left(\alpha(q^2)^{\frac{3}{2}}\right) \right]$$

Their first term is that of a free theory, consistent with asymptotic freedom. If we use only this first term of the free energy, we can derive

$$S = \frac{4}{3} \left[\frac{8\pi^2 V}{15} \left(1 + \frac{21}{32} n_{\mathrm{f}}\right) \right]^{\frac{1}{4}} E^{\frac{3}{4}}$$

which is the first term in our (3) with $d = 3$, $s = 16$ and $s^* = 12 n_{\mathrm{f}}$. However, there are corrections to the free energy coming from the fact that QCD is not free. These corrections that depend on q^2 (and therefore on the temperature) cannot simply be absorbed into the value of a. Unlike with Super Yang-Mills, there is no simple adjustment to (2) that would allow it to account for the coupling constant. One would have to already know that it is close to zero *a priori*.

The derivation will be done in three main steps. In the first step, our thermal partition function is calculated on $M \times \{\text{time}\}$ where M is a compact d-dimensional manifold. This manifold may or may not have a boundary. In the second step, the density of states is expressed as a Taylor series which is found by taking the inverse Laplace transform of the partition function (transitioning from a canonical to a microcanonical ensemble). In the last step, a more convenient expression is developed by finding the linear ODE satisfied by the Taylor series and employing the method of dominant balance. After the main result is derived, some aspects of the derivation are examined further. In particular, the specialization to one spatial dimension deserves its own discussion. There is also a generalization of the Cardy formula for arbitrary dimension proposed in 2000 by Erik Verlinde[6]. Determining the scope of its validity has received a large amount of attention[17–21]. The extent to which our results agree with this Cardy-Verlinde formula will be discussed at the end.

II. CALCULATING THE DENSITY OF STATES

A. The partition function

A field theory can be regarded as a collection of degrees of freedom labelled by a momentum $\mathbf{p}$ and possibly other quantum numbers labelling the spin and/or charge. When a particular degree of freedom having squared momentum p^2 is excited, the energy added to the system is $\sqrt{p^2+m^2}$ by the relativistic dispersion relation. In a bosonic theory, this excitation can be repeated infinitely many times, so if we restrict our attention to this degree of freedom, the single mode partition function is:

$$\begin{aligned} Z(p^2) &= 1 + e^{-\beta\sqrt{p^2+m^2}} + e^{-2\beta\sqrt{p^2+m^2}} + \dots \\ &= \left(1 - e^{-\beta\sqrt{p^2+m^2}}\right)^{-1} \end{aligned} \tag{4}$$

How many degrees of freedom have a squared momentum of p^2? This question depends on the manifold, M. Allowed momenta are eigenvalues of the Laplace-Beltrami operator. This is a map between the two function spaces $\Delta : C^2_{\mathrm{v}}(M) \to C^0(M)$ where the subscript v indicates Dirichlet boundary conditions. The eigenvalue equation is:

$$\Delta f = -\lambda f$$

where $\lambda = p^2$. Understanding the spectrum of this operator is related to Kac's problem; "Can you hear the shape of a drum?" Weyl's formula - one of the earliest positive answers to this question - will be immensely useful to us. Weyl's formula states that $\#(\lambda)$, the number of Dirichlet eigenvalues up to λ is given by:

$$\#(\lambda) = (2\pi)^{-d}\omega_d V \lambda^{\frac{d}{2}} + O\left(\lambda^{\frac{d}{2}-1}\right)$$

where V is the volume of the manifold and ω_d is the volume of a unit ball in $\mathbb{R}^d$ [22, 23]. The $O\left(\lambda^{\frac{d}{2}-1}\right)$ indicates that this asymptotic formula for the number of eigenvalues becomes increasingly accurate when λ is large. By differentiating this quantity, we can determine $g(\lambda)$ - the number of eigenvalues between λ and $\lambda + \mathrm{d}\lambda$.

$$\begin{aligned} g(\lambda) &= \frac{\mathrm{d}\#}{\mathrm{d}\lambda} \\ &\sim \frac{d}{2}(2\pi)^{-d}\omega_d V \lambda^{\frac{d}{2}-1} \end{aligned}$$

At this point, it is worth clearing up a potential source of confusion. While $g(\lambda)$ gives the degeneracy of the eigenvalue λ and λ is a way of labelling energy, $g(\lambda)$ is *not* the same as $\rho(E)$. In single particle problems, Weyl's formula is sometimes referred to as a formula for the density of states[24], but in this problem, we are interested in the density of states for the entire Fock space. We will therefore use the term *degeneracy* for $g(\lambda)$ and reserve the term *density of states* for $\rho(E)$. While this formula for $g(\lambda)$ gives us a degeneracy for the eigenvalue λ coming from geometrical considerations, we know that a field theory can introduce further degeneracy as part of its description. Therefore, if s is the number of internal states, there are really $sg(\lambda)$ ways of exciting a field to an eigenstate that has a squared momentum of λ. These excitations contribute a factor of

$$Z(\lambda)^{sg(\lambda)} = \left(1 - e^{-\beta\sqrt{\lambda+m^2}}\right)^{-sg(\lambda)}$$

to the partition function. The full partition function we seek is the product of $Z(\lambda)$ over all eigenvalues λ. Taking the log of a product turns it into a sum, so $\log Z = \sum_\lambda sg(\lambda)\log Z(\lambda)$. In the large volume limit, the spectrum of the Laplacian becomes continuous, meaning that our sum over λ becomes an integral.

$$\begin{aligned}
\log Z &= \int_0^\infty sg(\lambda)\log Z(\lambda)\mathrm{d}\lambda \\
&= -\frac{sd\omega_d V}{2(2\pi)^d}\int_0^\infty \log\left(1 - e^{-\beta\sqrt{\lambda+m^2}}\right)\lambda^{\frac{d}{2}-1}\mathrm{d}\lambda \\
&= \frac{sd\omega_d V}{2(2\pi)^d}\sum_{n=1}^\infty \int_0^\infty \frac{1}{n}e^{-n\beta\sqrt{\lambda+m^2}}\lambda^{\frac{d}{2}-1}\mathrm{d}\lambda \\
&= \frac{sd\omega_d V}{(2\pi)^d}\sum_{n=1}^\infty \int_0^\infty \frac{1}{n}e^{-n\beta\sqrt{p^2+m^2}}p^{d-1}\mathrm{d}p
\end{aligned}$$

In the second last step, we have Taylor expanded the logarithm and in the last step, we have used the fact that $\lambda = p^2$. To proceed further, we must make the field theory conformal by setting the mass to zero. This allows us to use the identity $\int_0^\infty e^{-cx}x^{d-1}\mathrm{d}x = \frac{(d-1)!}{c^d}$.

$$\begin{aligned}
\log Z &= \frac{sd!\omega_d V}{(2\pi\beta)^d}\sum_{n=1}^\infty \frac{1}{n^{d+1}} \\
&= \frac{sd!\omega_d\zeta(d+1)V}{(2\pi\beta)^d} \qquad (5)
\end{aligned}$$

The analysis so far has been applied to a bosonic theory, but very little changes when applying it to fermions. Each mode can have 0 or 1 excitations so instead of (4), we have:

$$Z(\lambda) = 1 + e^{-\beta\sqrt{\lambda}} \tag{6}$$

Proceeding to calculate (5) in the same way, we get:

$$\begin{aligned}
\log Z &= \int_0^\infty s g(\lambda) \log Z(\lambda) \mathrm{d}\lambda \\
&= \frac{sd\omega_d V}{2(2\pi)^d} \int_0^\infty \log\left(1 + e^{-\beta\sqrt{\lambda}}\right) \lambda^{\frac{d}{2}-1} \mathrm{d}\lambda \\
&= \frac{sd\omega_d V}{2(2\pi)^d} \sum_{n=1}^\infty \int_0^\infty \frac{(-1)^{n+1}}{n} e^{-n\beta\sqrt{\lambda}} \lambda^{\frac{d}{2}-1} \mathrm{d}\lambda \\
&= \frac{sd\omega_d V}{(2\pi)^d} \sum_{n=1}^\infty \int_0^\infty \frac{(-1)^{n+1}}{n} e^{-n\beta p} p^{d-1} \mathrm{d}p \\
&= \frac{sd!\omega_d V}{(2\pi\beta)^d} \sum_{n=1}^\infty \frac{(-1)^{n+1}}{n^{d+1}} \\
&= \frac{sd!\omega_d \zeta^*(d+1) V}{(2\pi\beta)^d}
\end{aligned} \tag{7}$$

where $\zeta^*(\sigma) = \left(1 - 2^{1-\sigma}\right)\zeta(\sigma)$ is the alternating zeta function. The functional form $Z = e^{\frac{a}{\beta^d}}$ will be essential in the next step where we link the partition function to the density of states.

B. From Laplace-Beltrami operators to Laplace transforms

We are now interested in finding $\rho(E)$ from $Z(\beta)$. By definition, the partition function is the Boltzmann factor $e^{-\beta E}$ summed (or integrated) over all states. Grouping the Boltzmann factors by energy makes the following formula apparent:

$$Z = \int_0^\infty \rho(E) e^{-\beta E} \mathrm{d}E$$

This is recognizable as the Laplace transform of the density of states so it is natural for us to consider the inverse Laplace transform of $e^{\frac{a}{\beta^d}}$. The following manipulations rely on a few straightforward identities involving hyperbolic functions.

$$\begin{aligned}
\rho(E) &= \frac{1}{2\pi i}\lim_{T\to\infty}\int_{\epsilon-iT}^{\epsilon+iT} e^{\beta E} e^{\frac{a}{\beta^d}}\mathrm{d}\beta \\
&= \frac{1}{2\pi i}\lim_{T\to\infty}\int_{\epsilon-iT}^{\epsilon+iT} e^{\beta E}\left[1+\frac{2}{\coth\left(\frac{a}{2\beta^d}\right)-1}\right]\mathrm{d}\beta \\
&= \frac{1}{2\pi}\int_{-\infty}^{\infty} e^{i\beta E}\mathrm{d}\beta + \frac{1}{\pi i}\lim_{T\to\infty}\int_{\epsilon-iT}^{\epsilon+iT} e^{\beta E}\frac{1}{\coth\left(\frac{a}{2\beta^d}\right)-1}\mathrm{d}\beta \\
&= \delta(E) + \frac{1}{\pi i}\lim_{T\to\infty}\int_{\epsilon-iT}^{\epsilon+iT} e^{\beta E}\frac{\tanh\left(\frac{a}{2\beta^d}\right)}{1-\tanh\left(\frac{a}{2\beta^d}\right)}\mathrm{d}\beta \\
&= \delta(E) + \frac{1}{\pi i}\lim_{T\to\infty}\int_{\epsilon-iT}^{\epsilon+iT} e^{\beta E}\frac{\tanh\left(\frac{a}{2\beta^d}\right)+\tanh^2\left(\frac{a}{2\beta^d}\right)}{1-\tanh^2\left(\frac{a}{2\beta^d}\right)}\mathrm{d}\beta \\
&= \delta(E) + \frac{1}{\pi i}\lim_{T\to\infty}\int_{\epsilon-iT}^{\epsilon+iT} e^{\beta E}\left[\frac{1}{2}\sinh\left(\frac{a}{2\beta^d}\right)+\sinh^2\left(\frac{a}{2\beta^d}\right)\right]\mathrm{d}\beta
\end{aligned}$$

To proceed further, we will Taylor expand $e^{\beta E}$. The interesting part of the density of states (the part without the delta function) now splits into two pieces.

$$\sum_{k=0}^{\infty}\frac{E^k}{k!}\lim_{T\to\infty}\left[\frac{1}{2\pi i}\int_{\epsilon-iT}^{\epsilon+iT}\beta^k\sinh\left(\frac{a}{\beta^d}\right)\mathrm{d}\beta+\frac{1}{\pi i}\int_{\epsilon-iT}^{\epsilon+iT}\beta^k\sinh^2\left(\frac{a}{2\beta^d}\right)\mathrm{d}\beta\right] \tag{8}$$

To evaluate the first integral, we will make the substitution $\alpha = \frac{1}{\beta}$.

$$\frac{1}{2\pi i}\lim_{T\to\infty}\int_{\frac{\epsilon-iT}{\epsilon^2+T^2}}^{\frac{\epsilon+iT}{\epsilon^2+T^2}}\frac{\sinh\left(a\alpha^d\right)}{\alpha^{k+2}}\mathrm{d}\alpha$$

The integrand has a pole at $\alpha = 0$ and the contour over which we are integrating is a vertical line, slightly to the right of this pole, with a vanishingly small height.



(a) Closed contour

(b) Open contour

FIG. 1: The integration to consider in the complex plane.

The figure on the right shows the desired contour, but fortunately, the integral over the contour shown on the left is equal. The leftward and rightward pieces of the contour cancel out while the downward piece is at infinity and does not contribute. By the residue theorem,

$$\begin{aligned}
\frac{1}{2\pi i}\lim_{T\to\infty}\int_{\frac{\epsilon-iT}{\epsilon^2+T^2}}^{\frac{\epsilon+iT}{\epsilon^2+T^2}}\frac{\sinh\left(a\alpha^d\right)}{\alpha^{k+2}}\mathrm{d}\alpha &= \operatorname{Res}\left(\frac{\sinh\left(a\alpha^d\right)}{\alpha^{k+2}};0\right)\\
&= \frac{1}{(k-d+1)!}\frac{\mathrm{d}^{k-d+1}}{\mathrm{d}\alpha^{k-d+1}}\left.\frac{\sinh\left(a\alpha^d\right)}{\alpha^d}\right|_{\alpha=0}\\
&= \frac{1}{(k-d+1)!}\frac{\mathrm{d}^{k-d+1}}{\mathrm{d}\alpha^{k-d+1}}\sum_{n=0}^{\infty}\left.\frac{a^{2n+1}\alpha^{2dn}}{(2n+1)!}\right|_{\alpha=0}\\
&= \begin{cases}\frac{a^{\frac{k+1}{d}}}{\left(\frac{k+1}{d}\right)!} & \frac{k+1}{d}\text{ odd}\\ 0 & \text{otherwise}\end{cases}
\end{aligned}$$

The second integral in (8) can be computed using the same substitution and the same contour. All that changes is the parity of the residue.

$$
\begin{aligned}
\frac{1}{\pi i}\lim_{T\to\infty}\int_{\frac{\epsilon-iT}{\epsilon^2+T^2}}^{\frac{\epsilon+iT}{\epsilon^2+T^2}}\frac{\sinh^2\left(\frac{a}{2}\alpha^d\right)}{\alpha^{k+2}}\mathrm{d}\alpha &= 2\mathrm{Res}\left(\frac{\sinh^2\left(\frac{a}{2}\alpha^d\right)}{\alpha^{k+2}};0\right) \\
&= \frac{2}{(k-2d+1)!}\frac{\mathrm{d}^{k-2d+1}}{\mathrm{d}\alpha^{k-2d+1}}\left.\frac{\sinh^2\left(\frac{a}{2}\alpha^d\right)}{\alpha^{2d}}\right|_{\alpha=0} \\
&= \frac{1}{(k-2d+1)!}\frac{\mathrm{d}^{k-2d+1}}{\mathrm{d}\alpha^{k-2d+1}}\left.\sum_{n=1}^{\infty}\frac{a^{2n}\alpha^{2d(n-1)}}{(2n)!}\right|_{\alpha=0} \\
&= \begin{cases} \frac{a^{\frac{k+1}{d}}}{\left(\frac{k+1}{d}\right)!} & \frac{k+1}{d}\text{ even} \\ 0 & \text{otherwise} \end{cases}
\end{aligned}
$$

We can now substitute both of these back into (8).

$$
\begin{aligned}
\rho(E) &= \delta(E)+\sum_{\frac{k+1}{d}\text{ odd}}\frac{E^k}{k!}\frac{a^{\frac{k+1}{d}}}{\left(\frac{k+1}{d}\right)!}+\sum_{\frac{k+1}{d}\text{ even}}\frac{E^k}{k!}\frac{a^{\frac{k+1}{d}}}{\left(\frac{k+1}{d}\right)!} \\
&= \delta(E)+\sum_{\frac{k+1}{d}=1}^{\infty}\frac{E^k}{k!}\frac{a^{\frac{k+1}{d}}}{\left(\frac{k+1}{d}\right)!} \\
&= \delta(E)+\sum_{j=1}^{\infty}\frac{a^jE^{dj-1}}{j!(dj-1)!} \qquad (9)
\end{aligned}
$$

When $d = 1$, this series can be expressed as a modified Bessel function but for other values of d this is surely something much less ubiquitous.

C. An asymptotic series from a Taylor series

In the previous step, the density of states we seek was expressed as a Taylor series, but not one with an elementary functional form. To better understand this Taylor series, we will define the function $f(x) = \sum_{m=1}^{\infty} b_m x^m$ where $b_m = \frac{1}{m!(dm-1)!}$. The recurrence relation satisfied by the b_m is $m(dm-1)(dm-2)\dots(dm-d)b_m = b_{m-1}$. Multiplying both sides by x^m and summing,

$$\sum_{m=1}^{\infty}\frac{1}{d}(md)\dots(md-d)b_m x^m - b_m x^{m+1} = 0$$

If x^m were x^{dm}, we could differentiate it $d+1$ times to turn it into $(md)\dots(md-d)x^{dm-d-1}$ and then multiply by x^{d+1} to get it into the form above. This suggests that our differential equation satisfied by f is:

$$\frac{1}{d}y^{d+1}\frac{\mathrm{d}^{d+1}}{\mathrm{d}y^{d+1}}\, f(y^d)\big|_{y=x^{\frac{1}{d}}} - xf(x) = 0$$
$$\frac{1}{d}x^{\frac{d}{d+1}}\frac{\mathrm{d}^{d+1}}{\left(\mathrm{d}x^{\frac{1}{d}}\right)^{d+1}} f(x) - xf(x) = 0 \tag{10}$$

For a fixed d, this equation can be expressed in a more standard form. The first three of them are:

d	
1	$x^2\frac{\mathrm{d}^2}{\mathrm{d}x^2}f(x) - xf(x) = 0$
2	$4x^3\frac{\mathrm{d}^3}{\mathrm{d}x^3}f(x) + 6x^2\frac{\mathrm{d}^2}{\mathrm{d}x^2}f(x) - xf(x) = 0$
3	$27x^4\frac{\mathrm{d}^4}{\mathrm{d}x^4}f(x) + 108x^3\frac{\mathrm{d}^3}{\mathrm{d}x^3}f(x) + 60x^2\frac{\mathrm{d}^2}{\mathrm{d}x^2}f(x) - xf(x) = 0$

(11)

It can be seen without much trouble that when a power of x appears beside a derivative, the exponent is equal to the order of the derivative. While it is rather technical to find the asymptotic series for $f(x)$, it can easily be converted into the asymptotic series for $\rho(E)$ because $\rho(E) = \frac{1}{E}f(aE^d)$.

Fact 1. *The coefficient on the highest order term in (10) is* d^d.

Proof. This is clear because $\frac{\mathrm{d}^{d+1}}{\mathrm{d}y^{d+1}}f(y^d) = \frac{\mathrm{d}^d}{\mathrm{d}y^d}\left[dy^{d-1}f'(y^d)\right]$. Every time one differentiates f, one gets another factor of dy^{d-1}. Doing this d more times yields $d^{d+1}y^{d^2-1}f^{(d+1)}(y^d)$. Letting $y = x^{\frac{1}{d}}$ and multiplying by $\frac{1}{d}x^{\frac{d+1}{d}}$, we get $d^dx^{d+1}f^{(d+1)}(x)$. $\square$

We will now set $f(x) = e^{S_0(x)}$ and observe that the highest order derivative gives us the term $d^dx^{d+1}\left(\frac{\mathrm{d}S_0}{\mathrm{d}x}\right)^{d+1}$ after we cancel $e^{S_0(x)}$. The other terms will all be of the form $\left(\frac{\mathrm{d}S_0}{\mathrm{d}x}\right)^{n_1}\dots\left(\frac{\mathrm{d}^jS_0}{\mathrm{d}x^j}\right)^{n_j}x^j$ where $n_1 + 2n_2 + \cdots + jn_j = j$ and $j \leq d+1$.

Fact 2. *Choosing* $S_0(x) \propto x^{\frac{1}{d+1}}$ *causes all such* $\left(\frac{\mathrm{d}S_0}{\mathrm{d}x}\right)^{n_1}\dots\left(\frac{\mathrm{d}^jS_0}{\mathrm{d}x^j}\right)^{n_j}x^j$ *to be* $o\left(x^{d+1}\left(\frac{\mathrm{d}S_0}{\mathrm{d}x}\right)^{d+1}\right)$.

Proof. Substituting our guess for $S_0(x)$ into the term above, we find that it is proportional to $\left(\frac{1}{x^{\frac{d}{d+1}}}\right)^{n_1}\dots\left(\frac{1}{x^{\frac{d}{d+1}+j-1}}\right)^{n_j}x^j = x^{j-n_1\left(\frac{d}{d+1}\right)-n_2\left(\frac{d}{d+1}+1\right)-\cdots-n_j\left(\frac{d}{d+1}+j-1\right)}$. We need to see if this is $o(x)$. Clearly it is because to maximize $j - n_1\left(\frac{d}{d+1}\right) - n_2\left(\frac{d}{d+1}+1\right) - \cdots - n_j\left(\frac{d}{d+1}+j-1\right)$, we only want to be subtracting the n_1 part, so we let all other $n_i = 0$. We now want to minimize $j\left(1-\frac{d}{d+1}\right)$ and the way to do this is to let $j = d+1$, giving us x. Anything less will give us something $o(x)$. □

With the knowledge that we can make all terms but $d^d x^{d+1}\left(\frac{\mathrm{d}S_0}{\mathrm{d}x}\right)^{d+1}$ negligible, our differential equation becomes:

$$\begin{aligned} d^d x^{d+1}\left(\frac{\mathrm{d}S_0}{\mathrm{d}x}\right)^{d+1} &= x \\ \frac{\mathrm{d}S_0}{\mathrm{d}x} &= \frac{1}{d^{\frac{d}{d+1}}} x^{\frac{1}{d+1}-1} \\ S_0(x) &= \frac{d+1}{d^{\frac{d}{d+1}}} x^{\frac{1}{d+1}} \end{aligned} \tag{12}$$

The solution to the differential equation when we drop all but one term, is precisely the solution that we have shown in Fact 2 to be consistent with the dropping of the terms in the first place. Thus, to zeroth order, the asymptotic behaviour of $f(x)$ must be $\log f(x) \sim \left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{1}{d+1}} x^{\frac{1}{d+1}}$. With a little more work, we can figure out what it is to first order.

Fact 3. *The coefficient on the second highest order derivative in (10) is $d^d \frac{d^2-1}{2}$.*

Proof. Again, starting with $\frac{\mathrm{d}^{d+1}}{\mathrm{d}y^{d+1}} f(y^d) = \frac{\mathrm{d}^d}{\mathrm{d}y^d}\left[d y^{d-1} f'(y^d)\right]$, we have to differentiate y^{d-1} once and differentiate f every other time in order to get an $f^{(d)}(y^d)$. If we wait until we have $\frac{\mathrm{d}}{\mathrm{d}y}\left[d^d y^{d(d-1)} f^{(d)}(y^d)\right]$, the last derivative gives us $d^d d(d-1) y^{d(d-1)-1} f^{(d)}(y^d)$. If we were to differentiate the factor of y earlier when it had $y^{r(d-1)}$, we would get $d^d r(d-1) y^{d(d-1)-1} f^{(d)}(y^d)$. Adding these up, we get $d^d(d-1)\left(\sum_{r=1}^{d} r\right) y^{d^2-d-1} f^{(d)}(y^d) = d^{d+1}\frac{d^2-1}{2} y^{d^2-d-1} f^{(d)}(y^d)$. Using $y = x^{\frac{1}{d}}$ and postmultiplying by $\frac{1}{d} x^{\frac{d+1}{d}}$ again, this becomes $d^d \frac{d^2-1}{2} x^d f^{(d)}(x)$. □

Since we have a solution for $S_0(x)$, we will now let $f(x) = e^{S_0(x)+S_1(x)}$. When this is inserted into (10), the highest power of x multiplying a single derivative of $S_1(x)$ will come from the first term in the expansion of

$$d^d x^{d+1}\left(\frac{1}{d+1}\frac{d+1}{d^{\frac{d}{d+1}}} x^{-\frac{d}{d+1}} + \frac{\mathrm{d}S_1}{\mathrm{d}x}\right)^{d+1} \tag{13}$$

and it will be $d\left(\frac{d}{d+1}\right)^{d-1}\left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{d}{d+1}} x^{d+1-\frac{d^2}{d+1}}\left(\frac{\mathrm{d}S_1}{\mathrm{d}x}\right)$. Also, by looking at the zeroth term in the expansion of (13), we see that an x exactly cancels the $-x$ present in the definition of (10).

Fact 4. *Choosing* $S_1(x) \propto \log x$ *makes* $x^{d+1-\frac{d^2}{d+1}}\left(\frac{\mathrm{d}S_1}{\mathrm{d}x}\right) = x^{\frac{d}{d+1}}$ *the dominant term.*

Proof. Recall that when $S_0(x)$ was the exponent, we had terms that looked like $\left(\frac{\mathrm{d}S_0}{\mathrm{d}x}\right)^{n_1} \dots \left(\frac{\mathrm{d}^j S_0}{\mathrm{d}x^j}\right)^{n_j} x^j$ where $n_1 + 2n_2 + \dots + jn_j = j$ and $j \leq d+1$. This time, we replace $S_0(x)$ with $x^{\frac{1}{d+1}} + S_1(x)$ so this product looks like $\left(x^{\frac{1}{d+1}-1} + \frac{\mathrm{d}S_1}{\mathrm{d}x}\right)^{n_1} \dots \left(x^{\frac{1}{d+1}-j} + \frac{\mathrm{d}^j S_1}{\mathrm{d}x^j}\right)^{n_j} x^j$. Terms in the expansion of this product look like:

$$\begin{aligned}
& x^{k_1\left(\frac{1}{d+1}-1\right)}\left(\frac{\mathrm{d}S_1}{\mathrm{d}x}\right)^{n_1-k_1} \dots x^{k_j\left(\frac{1}{d+1}-j\right)}\left(\frac{\mathrm{d}^j S_1}{\mathrm{d}x^j}\right)^{n_j-k_j} x^j \\
\propto\ & x^j x^{\frac{k_1+\dots+k_j}{d+1}-k_1-2k_2-\dots-jk_j-(n_1-k_1)-2(n_2-k_2)-\dots-j(n_j-k_j)} \\
=\ & x^{\frac{k_1+\dots+k_j}{d+1}}
\end{aligned}$$

To maximize the exponent, we must turn it into $\frac{n_1+\dots+n_j}{d+1}$ and the way to maximize this is to let $n_1 = j = d+1$. However, we have aleady mentioned that the x term should vanish so the next highest power of x we can get is $x^{\frac{d}{d+1}}$. We realize this either with $n_1 = j = d$ or with $j = d+1$ and $k_1 = n_1 - 1$. Everything else is $o\left(x^{\frac{d}{d+1}}\right)$. ☐

This allows us to drop most of the terms in (10) just like before. Specifically, we will drop all powers of x, lower than $x^{\frac{d}{d+1}}$ and find that this is consistent with having $S_1(x) \propto \log x$. The next two facts are devoted to solving for the coefficients that appear beside $x^{\frac{d}{d+1}}$.

Fact 5. $\frac{\mathrm{d}^n}{\mathrm{d}x^n} e^{rx^m}$ *can be written as* $x^{-n} e^{rx^m} \sum_{j=0}^{n} a_j (rx^m)^j$.

Proof. We are dealing with $\frac{\mathrm{d}^{n-1}}{\mathrm{d}x^{n-1}}\left(mrx^{m-1}e^{rx^m}\right)$. If we differentiate the exponent k times, we get $m^k r^k x^{k(m-1)} e^{rx^m}$. If we then differentiate the power of x $n-k$ times, our result is proportional to the exponential times $r^k x^{k(m-1)-(n-k)=x^{-n}(rx^m)^k}$. ☐

In the situations where we need to apply this fact, $m = \frac{1}{d+1}$ and $r = \frac{d+1}{d^{\frac{d}{d+1}}}$. One way to get an $x^{\frac{d}{d+1}}$ term is to let $n = j = d$. In this case, the coefficient we find is $r^d m^d =$

$\left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{d}{d+1}} \frac{1}{(d+1)^d}$. We must remember that this is multiplied by the coefficient on the d^{th} derivative found in Fact 3.

Fact 6. *The other route to an* $x^{\frac{d}{d+1}}$ *term - choosing* $n = d+1$ *and* $j = d$ *- yields a coefficient of* $\frac{-d^2}{2}\frac{1}{d^{\frac{d^2}{d+1}}}$.

Proof. We have $\frac{\mathrm{d}^d}{\mathrm{d}x^d}\left(mrx^{m-1}e^{rx^m}\right)$ and we have to differentiate the exponential all but one time. Waiting until $\frac{\mathrm{d}}{\mathrm{d}x}\left(m^d r^d x^{d(m-1)}e^{rx^m}\right)$, this differentiates to $d(m-1)m^d r^d x^{dm-d-1}e^{rx^m}$. Had we done this for $x^{s(m-1)}$, we would have had $s(m-1)m^d r^d x^{dm-d-1}$ times the exponential. We can therefore add these up to get $\frac{-d}{d+1}\left(\frac{1}{d+1}\right)^d\left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{d}{d+1}}\sum_{s=1}^{d} s = \frac{-d^2}{2}\frac{1}{d^{\frac{d^2}{d+1}}}$. ☐

We must also remember that this is multiplied by the coefficient on the $(d+1)^{\text{st}}$ derivative found in Fact 1. The following equation is the result of keeping only the terms found to be significant in Fact 4 complete with prefactors. The left hand side uses the coefficient observed from (13) and the right hand side uses the two coefficients found above corresponding to the two ways of constructing $x^{\frac{d}{d+1}}$.

$$\begin{aligned}
d\left(\frac{d}{d+1}\right)^{d-1}\left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{d}{d+1}} x^{1+\frac{d}{d+1}}\left(\frac{\mathrm{d}S_1}{\mathrm{d}x}\right) &= \left[d^d\frac{d^2}{2}\frac{1}{d^{\frac{d^2}{d+1}}} - d^d\frac{d^2-1}{2}\frac{1}{d^{\frac{d^2}{d+1}}}\right]x^{\frac{d}{d+1}} \\
\frac{\mathrm{d}S_1}{\mathrm{d}x} &= \frac{1}{x}\left[\frac{d^2}{2(d+1)} - \frac{d-1}{2}\right] \\
S_1(x) &= \frac{1}{2(d+1)}\log x + \log C_d \qquad (14)
\end{aligned}$$

Finally substituting (12) and (14) into $f(x) = e^{S_0(x)+S_1(x)}$, we obtain the result:

$$\begin{aligned}
f(x) &\sim C_d x^{\frac{1}{2(d+1)}} e^{\left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{1}{d+1}} x^{\frac{1}{d+1}}} \\
\rho(E) &\sim C_d a^{\frac{1}{2(d+1)}} E^{\frac{-d-2}{2(d+1)}} e^{\left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{1}{d+1}} a^{\frac{1}{d+1}} E^{\frac{d}{d+1}}} \qquad (15)
\end{aligned}$$

A remaining point to address is the integration constant C_d. We used the method of dominant balance to find the asymptotic behaviour of f, a function defined by a Taylor series. However, what we really did was find the asymptotic behaviour of any function that solves the same differential equation as f. This equation is linear, so there is no way that such an analysis can tell us the constant by which the expression (15) should be multiplied. The integration constant can be fixed if we substitute the same large number into f's Taylor series and f's asymptotic series and demand that these expressions be equal. This has been done for several values of d in the plot. Looking back to (1), it is no coincidence that our computed value of C_1 is very close to $\frac{1}{2\sqrt{\pi}}$.



FIG. 2: The overall constant appearing in the asymptotic series for the density of states as a function of dimension.

It is also worth noting that this method could in principle be continued to find additional terms in the asymptotic series for $\rho(E)$. The only differential equation (10) we could write down for a general d had coefficients that were not very explicit. This fact would make the higher order terms progressively more difficult to compute. However, for a fixed d, there is no reason to stop at the $f(x) = e^{S_0(x)+S_1(x)}$ ansatz. For $d = 1$, it is a simple exercise to find the more accurate asymptotic expansion

$$\begin{aligned} f(x) &\sim C_1 x^{\frac{1}{4}} e^{2\sqrt{x}-\frac{3}{16\sqrt{x}}} \\ \rho(E) &\sim C_1 a^{\frac{1}{4}} E^{-\frac{3}{4}} e^{2\sqrt{aE}-\frac{3}{16\sqrt{aE}}} \end{aligned} \tag{16}$$

which comes from considering $f(x) = e^{S_0(x)+S_1(x)+S_2(x)}$ when $S_0(x)$ and $S_1(x)$ are already known.

D. Limitations

Going back to our calculation of the partition function, a formula we used was:

$$\log Z = \frac{s^{(*)} d\omega_d V}{(2\pi)^d} \sum_{n=1}^{\infty} \int_0^{\infty} \frac{(\pm 1)^{n+1}}{n} e^{-n\beta p} p^{d-1} \mathrm{d}p$$

The approximation of a continuous momentum spectrum is only useful if the large momenta contribute significantly to Z. This contribution vanishes as we raise the value of β so our formula for the partition function is only valid at high temperatures. Another way to say this is that we want the correction to Weyl's formula to be small. Weyl's formula for $\#(p^2)$ is roughly a degree d polynomial in p. The term that we have been using is proportional to the volume of the manifold but there is a second term, found by Ivrii[22], that is proportional to the area of its boundary ∂V.

$$\#(\lambda) = (2\pi)^{-d} \omega_d V \lambda^{\frac{d}{2}} + \frac{1}{4} (2\pi)^{-(d-1)} \omega_{d-1} \partial V \lambda^{\frac{d}{2}-1} + o\left(\lambda^{\frac{d}{2}-1}\right)$$

This would give the log of the partition function a $\frac{\partial V}{\beta^{d-1}}$ term in addition to the $\frac{V}{\beta^d}$ that we found. Making this extra term small requires that the temperature be larger than the inverse length scale $\frac{\partial V}{V}$.

This high temperature constraint on the partition function translates to a high energy constraint on the density of states. It is easy to see this if we use the fact that the peak energy in a canonical ensemble goes as VT^{d+1}. Another justification comes from the formula for the density of states. βE must be on the order of 1 if we want the "incorrect parts" of the partition function to contribute negligibly to the inverse Laplace transform. If $E \gg \frac{\partial V}{V}$ were not satisfied, the $\rho(E) \propto e^{E^{\frac{d}{d+1}}}$ result that we found would in fact be a convolution of $e^{E^{\frac{d}{d+1}}}$, $e^{E^{\frac{d-1}{d}}}$, $e^{E^{\frac{d-2}{d-1}}}$, *etc* and the error arising from not computing this convolution would dominate the effect of the logarithmic correction.

One can imagine a procedure which "squishes" the manifold such that ∂V increases but V

stays the same. This will raise the energy required to make sense of our results, but as long as ∂V stays finite, such an energy will still exist. A different problem occurs if we take ∂V to zero. $E \gg \frac{\partial V}{V}$ is only the right condition if the dominant correction to Weyl's formula is proportional to ∂V. For a boundaryless manifold (such as a sphere), $E > 0$ is not sufficient to make our main result correct. When we discuss the Cardy-Verlinde formula, we will show that correct condition for a sphere of radius r is $E \gg r^{-1}$.

Knowing this, it is perhaps not useful that we used the method of dominant balance a third time to derive (16). We can Taylor expand it to get:

$$\begin{aligned}\rho(E) &\sim C_1 a^{\frac{1}{4}} E^{-\frac{3}{4}} e^{2\sqrt{aE}-\frac{3}{16\sqrt{aE}}} \\ &= C_1 a^{\frac{1}{4}} E^{-\frac{3}{4}} e^{2\sqrt{aE}} \left(1 - O\left(\frac{1}{\sqrt{rE}}\right)\right)\end{aligned}$$

In deriving this, we used the fact that $Er \gg 1$. However, the Taylor expansion tells us that the only term we derived by going beyond our main result, is one that we already assumed to be zero. By comparison, the additional assumptions required to apply our result to an asymptotically free theory, will not lead to their own inconsistencies. If $E \gg \Lambda_{\text{QCD}}$, we still only need to make sure that $Er \gg 1$. The formula for $\rho(E)$ will not yield any terms proportional to $\frac{1}{E\Lambda_{\text{QCD}}^{-1}}$.

III. DISCUSSION

A. Other approximations

The calculation of (15) was split into three steps but one may ask if all of these steps are really needed. Most of the progress is made in the first step. Once the formula $Z = e^{\frac{a}{\beta^d}}$ is derived and the constant a is known, there are two simple procedures that can be carried out to get an approximate density of states. Unsurprisingly, they give the same result.

The first procedure is to calculate the entropy in terms of temperature using

$$S = -\frac{\partial F}{\partial T} = \frac{\partial}{\partial T}(T \log Z) \tag{17}$$

One can then solve for the energy in terms of temperature using $E = -\frac{\partial \log Z}{\partial \beta}$. This is really the expected energy of the canonical ensemble, but energy fluctuations away from this are surpressed in the thermodynamic limit. Inverting this, one can insert temperature into (17) to find $S(E)$.

The second procedure is the one used by Cardy[7]. The power law $S = \tilde{C} E^{\frac{d}{d+1}}$ for some prefactor $\tilde{C}$ is known by conformal invariance. The definition of temperature states that $\frac{\partial S}{\partial E}|_{E_0} = \beta$ which can be used to solve for the expected energy E_0 in terms of β and $\tilde{C}$. In the saddle point approximation, $Z = e^{S(E_0) - \beta E_0}$. Demanding that this be equal to $e^{\frac{a}{\beta^d}}$ allows one to solve for $\tilde{C}$ and thereby solve for $S(E)$.

What we get from both these analyses is that:

$$S = \left(\frac{(d+1)^{d+1}}{d^d} \right)^{\frac{1}{d+1}} a^{\frac{1}{d+1}} E^{\frac{d}{d+1}}$$

If we exponentiate this, we get the zeroth term in the asymptotic series for $\rho(E)$ - our result (15) without the logarithmic correction. The role of the logarithmic correction can be understood from dimensional analysis. Boltzmann's formula states that $S = \log(\Omega(E))$ where $\Omega(E)$ is the exact number of states that have energy E. Another way to say this is that $\Omega(E)$ gives the number of states between E and $E + \mathrm{d}E$. However, the quantity $\Omega(E)$ is not very useful for the techniques that approximate the spectrum of a thermodynamic system as being continuous for large energies. What we seek instead, the *density* of states, gives the the number of states in $[E, E + \mathrm{d}E)$ as $\rho(E)\mathrm{d}E$ which requires it to have units of inverse energy. It is essentially $\Omega(E)$ times an inverse energy prefactor. In some sense, the second and third steps in our main calculation exist to solve for this prefactor as $a^{\frac{1}{2(d+1)}} E^{\frac{-d-2}{2(d+1)}}$.

Even though we focused on finding the microcanonical density of states, our analysis is similar in spirit to two studies of how the entropy fluctuates in a canonical ensemble[25, 26]. These authors also express the density of states as the inverse Laplace transform of the partition function. Rather than inserting a particular form for the partition function, they leave it general in order to obtain the entropy's deviation from $S(\beta_0)$ in terms of $S(\beta_0)$. The saddle point approximation is used to evaluate this inverse Laplace transform so that it has an exponent of $\log Z(\beta_0) - \beta_0 E$. Both papers agree that the fluctuation is:

$$\begin{aligned} S &= S_0 - \frac{1}{2}\log C_V \\ C_V &= \beta^2 \frac{\partial^2 \log Z}{\partial \beta^2}\Big|_{\beta_0} \end{aligned} \tag{18}$$

However, they use different methods to calculate C_V and [25] makes note of a conflict: [25] finds $S = S(\beta_0) - \frac{1}{2}\log S(\beta_0)$ while [26] finds $S = S(\beta_0) - \frac{3}{2}\log S(\beta_0)$. Since we have an explicit form of the partition function, we can see which result is reproduced in a free theory. The equilibrium inverse temperature we find from $Z = e^{\frac{a}{\beta^d}}$ is $\beta_0 = \left(\frac{da}{E}\right)^{\frac{1}{d+1}}$. By differentiating $\log Z$ twice and using this temperature, we find that:

$$C_V = (d+1)(da)^{\frac{1}{d+1}} E^{\frac{d}{d+1}} \propto E^{\frac{d}{d+1}}$$

Since this is proportional to $S(\beta_0)$ and not $S(\beta_0)^3$, entropy fluctuations in this setting appear to be in favour of [25].

B. Relation to CFT_2

The real strength of Cardy's approach lies in a duality between the low and high temperature limits of a CFT partition function.

The partition function for a CFT on a torus is usually expressed in terms of the *modular parameter* τ. A parallelogram is defined by two linearly independent vectors, and a torus is defined by a parallelogram if we identify the edges. If we express the two vectors as complex numbers, τ is defined to be their ratio. This means that τ is dimensionless. A generalized definition of the partition function is

$$Z = \mathrm{Tr}\left[e^{-V(H\,\mathrm{Im}(\tau) - iP\,\mathrm{Re}(\tau))}\right] \tag{19}$$

where H and P are the energy and momentum operators respectively and V is the spatial volume (one of the periods of the torus). This becomes the regular partition function from statistical mechanics if $\tau = \frac{i\beta}{V}$. Unitary conformal field theories possess the property of invariance under the modular group $\mathrm{PSL}(2,\mathbb{Z})$. Modular transformations take the form

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

where a, b, c and d are integers satisfying $ad - bc = 1$. They are generated by two transformations:

$$S : \tau \mapsto -\frac{1}{\tau} \qquad T : \tau \mapsto \tau + 1$$

To discuss modular invariance in the context of our result, we should consider what the low temperature limit of our partition function should be. At high values of β, all Boltzmann factors other than the one coming from the ground state are exponentially surpressed. The partition function is therefore dominated by the energy of the vacuum which is $-\frac{2\pi}{V}\frac{c+\tilde{c}}{24}$. Here, c and $\tilde{c}$ are the left and right central charges. Since they are equal in the theories we consider, we will call this $-\frac{\pi c}{6V}$. A proper partition function should therefore satisfy:

$$Z \to e^{\frac{\beta\pi c}{6V}} \text{ as } \beta \to \infty \tag{20}$$

Clearly, the partition function we wrote down earlier does not have this limit because we assumed high temperatures in its construction. However, this limit is all that we need to know to derive the high temperature limit as well. Inverse temperature and time are related by a Wick rotation so a conformal field theory in one spatial dimension can be regarded as a Euclidean CFT on a torus.



FIG. 3: A CFT in volume V at inverse temperature β and a CFT in volume V at inverse temperature $\frac{V^2}{\beta}$ are related by a conformal transformation.

Scaling both lengths by $\frac{V}{\beta}$ turns a V by β torus into a $\frac{V^2}{\beta}$ by V torus. This scaling is a conformal transformation and a Euclidean theory is agnostic to whether the first or the second co-ordinate carries temperature information. This tells us that we must have:

$$Z(\beta) = Z\left(\frac{V^2}{\beta}\right)$$

This provides the desired link between the high and low temperature partition functions:

$$Z \to e^{\frac{V\pi c}{6\beta}} \text{ as } \beta \to 0 \tag{21}$$

The low temperature limit (20) holds in any number of dimensions with any type of interaction. The high temperature limit (21) holds with interactions as well but only in two dimensions. The reason for this is clear from the derivation - the number of dimensions with and without temperature information have to be the same. The figure shows the common idiom $\beta \mapsto \frac{4\pi^2}{\beta}$. With $\tau = \frac{i\beta}{V}$, this is equivalent to switching β and V. In other words, this is an S transformation. In many cases, invariance of the partition function under S is simply called modular invariance[8, 27] because the effect of the T transformation is trivial.

Naturally the partition function given by (5) has the form of (21). Substituting $d = 1$, we have:

$$\begin{aligned} \log Z &= \frac{s\zeta(2)V}{\pi\beta} \\ &= \frac{V\pi s}{6\beta} \end{aligned}$$

Even though our derivation required no knowledge of the Casimir effect, we confirm that the central charge of a bosonic field is equal to the number of internal states[27]. This also holds for Dirac fermions which have $c = 1$[28]. The number of spin states is $s = 2$ and the required factor of $\frac{1}{2}$ comes from the zeta function identity $\zeta^*(\sigma) = \left(1 - 2^{1-\sigma}\right)\zeta(\sigma)$. It is therefore clear that if we choose $d = 1$, our formula for the partition function - and therefore the density of states - becomes valid for an interacting theory as well. As stated in the main text, the choice $d = 1$ also renders the differential equation exactly solvable so there is no need for asymptotic analysis. The Taylor series (9) becomes the Taylor series of a modified Bessel function giving the result of [8].

C. The Cardy-Verlinde formula

When the temperature is large compared to the inverse length scale, what we found to be a negligible effect of topology on the partition function is not surprising. In [29], the authors state that this behaviour is fully expected. They also give an example appearing in the AdS/CFT correspondence[1] which shows that the assertion may fail at low temperatures. As a consequence of Weyl's formula, we have seen that the leading term in the free energy is indeed independent of the manifold. However, the *sub-leading* behaviour need not be. The sub-leading free energy on a manifold with a boundary will look much different from the sub-leading free energy on a manifold without a boundary. This is why our final discussion will specialize to a sphere of radius r. As we will see, specializing to a manifold with a boundary instead would cause the Cardy-Verlinde formula to break down.

The form of the Cardy formula that we have mentioned multiple times is:

$$\begin{aligned} S &= 2\pi r\sqrt{\frac{c+\tilde{c}}{12r}\left(2E-\frac{c+\tilde{c}}{12r}\right)} \\ &\approx 2\pi\sqrt{\frac{crE}{3}} \end{aligned}$$

where r is the radius of the circle on which the CFT is defined. On a d-sphere, the Cardy-Verlinde formula[6] looks very similar:

$$S = \frac{2\pi r}{\sqrt{AB}}\sqrt{E_{\mathrm{C}}(2E-E_{\mathrm{C}})} \tag{22}$$

An entropy proportional to $\sqrt{E}$ rather than $E^{\frac{d}{d+1}}$ would obviously be not be correct so some care has to be taken in how the Casimir energy E_{C} is defined. Verlinde arrives at the formula using the relations:

$$\begin{aligned} E &= E_{\mathrm{E}} + \frac{1}{2}E_{\mathrm{C}} \\ E_{\mathrm{E}} &= \frac{A}{4\pi r}S^{1+\frac{1}{d}} \\ E_{\mathrm{C}} &= \frac{B}{2\pi r}S^{1-\frac{1}{d}} \end{aligned}$$

E_{E} is the *extensive* part of the energy, something that we have been careless enough to have simply called the energy all this time. Its dependence on $S^{1+\frac{1}{d}}$ should be familiar by this point. The non-trivial claim is that $E - E_{\mathrm{E}} \propto S^{1-\frac{1}{d}}$.

Taking $S \propto VT^d$, it suffices to show that $E_{\mathrm{C}} \propto T^{d-1}$. For $d > 1$, this contribution vanishes at zero temperature. It is therefore completely unrelated to the vaccum energies that are typically calculated using regularization techniques. Rather, in a free theory, it is related to the finite size effects in Weyl's formula. We have seen that the first term in Weyl's formula contributes VT^{d+1} to the free energy (the extensive part), while the second term contributes

$\partial V T^d$. This means that in order for the terms in the free energy to jump from T^{d+1} to T^{d-1}, the manifold must be *boundaryless.*

The Cardy-Verlinde formula has been most successful at describing strongly coupled theories with AdS duals. The main work studying its use in a free theory is a 2001 paper by Kutasov and Larsen[17]. They compute exact partition functions without passing to a continuum limit for free theories on $\mathbb{S}^1 \times \mathbb{S}^3$ and $\mathbb{S}^1 \times \mathbb{S}^5$ (some of their results can be checked in [5]). Most of their results show the desired T^{d-1} scaling and this is because they use a shape with an empty boundary as their spatial manifold. By examining spherical harmonics, we can perform a calculation in the continuum limit that agrees with the results of [17] except for the $d = 3$ scalar with conformal coupling.

The d-dimensional spherical harmonics obey the eigenvalue equation

$$r^2 \Delta Y_{l_1,\dots,l_d} = -l_d(l_d + d - 1) Y_{l_1,\dots,l_d}$$

and have integer indices that satisfy $0 \leq |l_1| \leq l_2 \leq \cdots \leq l_d$[30]. If the condition were $l_1 \leq \cdots \leq l_d$, we could use the formula for l_d-simplex numbers to solve for the degeneracy of each eigenvalue as $\binom{l_d+d-1}{d-1}$. However, l_1 is allowed to be negative. Multiplying by 2 and subtracting the number of $l_2 \leq \cdots \leq l_d$ choices so as not to double count $l_1 = 0$, we find a degeneracy given by:

$$g(l_d) = 2\binom{l_d + d - 1}{d - 1} - \binom{l_d + d - 2}{d - 2} = \frac{2l_d + d - 1}{d - 1}\binom{l_d + d - 2}{d - 2}$$

Using the eigenvalue equation and the degeneracy, we may write:

$$\begin{aligned}
\frac{2l_d + d - 1}{d - 1}\binom{l_d + d - 2}{d - 2}\mathrm{d}l_d &= \frac{\sqrt{(d-1)^2 + 4r^2p^2}}{d - 1}\binom{l_d + d - 2}{d - 2}\mathrm{d}l_d \\
&= \frac{\sqrt{(d-1)^2 + 4r^2p^2}}{d - 1}\binom{l_d + d - 2}{d - 2}\frac{2r^2p}{\sqrt{(d-1)^2 + 4r^2p^2}}\mathrm{d}p \\
&= \frac{2r^2p}{d - 1}\binom{l_d + d - 2}{d - 2}\mathrm{d}p
\end{aligned}$$

Carrying out the same procedure as before, we have:

$$\begin{aligned}
\log Z &= -\frac{2s}{d-1}\int_0^\infty r^2 p\binom{l_d+d-2}{d-2}\log\left(1-e^{-\beta p}\right)\mathrm{d}p \\
&= -\frac{2s}{(d-1)!}\int_0^\infty r^2(l_d+d-2)(l_d+d-3)\ldots(l_d+1)\log\left(1-e^{-\beta p}\right)p\mathrm{d}p \\
&= -\frac{2s}{(d-1)!}\int_0^\infty \frac{r^2}{2^{d-2}}(2l_d+2d-4)(2l_d+2d-6)\ldots(2l_d+2)\log\left(1-e^{-\beta p}\right)p\mathrm{d}p \\
&= -\frac{2s}{(d-1)!}\int_0^\infty \frac{r^2}{2^{d-2}}(\sqrt{4r^2p^2+(d-1)^2}+d-3)(\sqrt{4r^2p^2+(d-1)^2}+d-5) \\
&\quad \ldots(\sqrt{4r^2p^2+(d-1)^2}+3-d)\log\left(1-e^{-\beta p}\right)p\mathrm{d}p
\end{aligned}$$

Notice that the product in the integrand is already factored as a difference of squares. If $d=2$, it is the empty product. If d is an even number greater than 2, it is a product of $(4r^2p^2+(d-1)^2+(d-3)^2)$, $(4r^2p^2+(d-1)^2+(d-5)^2)$, *etc.* If d is odd, there is an extra $\sqrt{4r^2p^2+(d-1)^2}$ left over which we Taylor expand as $2rp+\frac{(d-1)^2}{4rp}$. Collecting all the highest powers of r in the integrand, we get r^dp^{d-1}. We would now like to find the term with the next highest power of r. In the even case, it is:

$$\frac{1}{4}r^{d-2}p^{d-3}\left[\left(\frac{d}{2}-1\right)(d-1)^2-\sum_{k=0}^{\frac{d-4}{2}}(2k+1)^2\right]=\frac{1}{12}r^{d-2}p^{d-3}(d^3-4d^2-11d-6)$$

In the odd case, it is:

$$\frac{1}{4}r^{d-2}p^{d-3}\left[\left(\frac{d}{2}-1\right)(d-1)^2-\sum_{k=0}^{\frac{d-3}{2}}(2k)^2\right]=\frac{1}{12}r^{d-2}p^{d-3}d(d^2-3d+11)$$

We can therefore evaluate

$$\begin{aligned}
\log Z &= -\frac{2s}{(d-1)!}\int_0^\infty (r^dp^{d-1}+\frac{1}{12}h(d)r^{d-2}p^{d-3})\log\left(1-e^{-\beta p}\right)\mathrm{d}p \\
&= \frac{2dsr^d\zeta(d+1)}{\beta^d}+\frac{sh(d)r^{d-2}\zeta(d-1)}{6(d-1)(d-2)\beta^{d-2}}
\end{aligned}$$

where

$$h(d)=\begin{cases}0 & d=2\\ d^3-4d^2-11d-6 & d\ \text{even}\\ d^3-2d^2+11d & d\ \text{odd}\end{cases} \tag{23}$$

Again the zeta function is the only thing that would change if we had done the calculation for fermions. Notice that 6 is a root of $h(d)$. We now have the extensive and sub-extensive parts of the free energy and the entropy:

$$\begin{aligned}
F &= -2d[s\zeta(d+1)+s^*\zeta^*(d+1)]r^dT^{d+1} - \frac{h(d)[s\zeta(d-1)+s^*\zeta^*(d-1)]}{6(d-1)(d-2)}r^{d-2}T^{d-1} \\
S &= 2d(d+1)[s\zeta(d+1)+s^*\zeta^*(d+1)]r^dT^d + \frac{h(d)[s\zeta(d-1)+s^*\zeta^*(d-1)]}{6(d-2)}r^{d-2}T^{d-2}
\end{aligned}$$

Defining the Casimir energy as $E + dF$ (the deviation from the Euler relation [6]), it is straightforward to compute the energies:

$$\begin{aligned}
E_{\mathrm{E}} &= 2d^2[s\zeta(d+1)+s^*\zeta^*(d+1)]r^dT^{d+1} + \frac{h(d)(d-3)[s\zeta(d-1)+s^*\zeta^*(d-1)]}{6(d-1)(d-2)}r^{d-2}T^{d-1} \\
E_{\mathrm{C}} &= \frac{h(d)[s\zeta(d-1)+s^*\zeta^*(d-1)]}{3(d-1)(d-2)}r^{d-2}T^{d-1}
\end{aligned}$$

Strictly speaking, neither is a pure power of S. However, if we only require the leading terms to be a power of the leading term of S, the coefficients we seek are:

$$\begin{aligned}
\frac{A}{4\pi r} &= \frac{d}{r\left(2d[s\zeta(d+1)+s^*\zeta^*(d+1)]\right)^{\frac{1}{d}}(d+1)^{1+\frac{1}{d}}} \\
\frac{B}{2\pi r} &= \frac{h(d)[s\zeta(d-1)+s^*\zeta^*(d-1)]}{3(d-1)(d-2)r}\left(2d(d+1)[s\zeta(d+1)+s^*\zeta^*(d+1)]\right)^{\frac{1}{d}-1}
\end{aligned}$$

Substituting these into (22), the corresponding Cardy-Verlinde formula is:

$$\begin{aligned}
S &= (d+1)\sqrt{\frac{3(d-1)(d-2)}{h(d)}\frac{s\zeta(d+1)+s^*\zeta^*(d+1)}{s\zeta(d-1)+s^*\zeta^*(d-1)}}r\sqrt{E_{\mathrm{C}}(2E-E_{\mathrm{C}})} \\
&= \left(\frac{(d+1)^{d+1}}{d^d}\right)^{\frac{1}{d+1}}\left(2d[s\zeta(d+1)+s^*\zeta^*(d+1)]\right)^{\frac{1}{d+1}} r^{\frac{d}{d+1}} E_{\mathrm{E}}^{\frac{d}{d+1}}
\end{aligned}$$

The holographic theories studied by Verlinde have $AB = d^2$. When Kutasov and Larsen say that the Cardy-Verlinde formula does not hold in a free theory, they are talking about this more restrictive type of Cardy-Verlinde formula. The formula above holds on any sphere that is not a 2-sphere or a 6-sphere and as noted in [17], the overall coefficient depends on the matter content of the theory. Extending this formula to a general boundaryless manifold would require us to know the third term in Weyl's formula.

IV. ACKNOWLEDGEMENTS

This work was supported in part by the Natural Sciences and Engineering Research Council of Canada. It also benefited from helpful comments given by Mark Van Raamsdonk, Michael McDermott and Jared Stang.

- [1] J. Maldacena. The large-N limit of superconformal field theories and supergravity. *International Journal of Theoretical Physics*, 38:1113–1133, 1998. arXiv:9711.200.
- [2] R. Rattazzi; S. Rychkov; A. Vichi. Central charge bounds in 4d conformal field theory. *Physical Review D*, 83, 2011. arXiv:1009.2725.
- [3] D. Z. Freedman; S. D. Mathur; A. Matusis; L. Rastelli. Correlation functions in the CFT(d)/AdS(d+1) correspondence. *Nuclear Physics B*, 546:96–118, 1999. arXiv:9804.058.
- [4] S. Ferrara; C. Fronsdal. Conformal fields in higher dimensions. 1998. arXiv:9804.058.
- [5] O. Aharony; D. Marolf; M. Rangamani. Conformal field theories in anti-de Sitter space. *Journal of High Energy Physics*, 2, 2011. arXiv:1011.6144.
- [6] E. Verlinde. On the holographic principle in a radiation dominated universe. 2000. arXiv:0008140.
- [7] J. L. Cardy. Operator content of two-dimensional conformally invariant theories. *Nuclear Physics B*, 270:186–204, 1986.
- [8] F. Loran; M. M. Sheikh-Jabbari; M. Vincon. Beyond logarithmic corrections to Cardy formula. *Journal of High Energy Physics*, 110, 2011. arXiv:1010.3561.
- [9] F. Brauer. On the asymptotic behaviour of Bessel functions. *American Mathematical Monthly*, 70(9):954–957, 1963.
- [10] S. S. Gubser; I. R. Klebanov; A. A. Tseytlin. Coupling constant dependence in the thermodynamics of N = 4 supersymmetric Yang-Mills theory. *Nuclear Physics B*, 534:202–222, 1998. arXiv:9805.156.
- [11] C. Kim; S. J. Rey. Thermodynamics of large-N super Yang-Mills theory and AdS/CFT correspondence. *Nuclear Physics B*, 564, 2000. arXiv:9905.205.

[12] A. Fotopoulos; T. R. Taylor. Remarks on two-loop free energy in N=4 supersymmetric Yang-Mills theory at finite temperature. *Physical Review D*, 59, 1999. arXiv:9811.224.

[13] R. G. Cai; N. Ohta. Thermodynamics of large N noncommutative super Yang-Mills theory. *Physical Review D*, 61, 2000. arXiv:9910.092.

[14] D. J. Gross; F. Wilczek. Ultraviolet behaviour of non-abelian gauge theories. *Physical Review Letters*, 30:1343–1346, 1973.

[15] H. D. Politzer. Reliable perturbative results for strong interactions? *Physical Review Letters*, 30:1346–1349, 1973.

[16] E. Braaten; A. Nieto. Free energy of QCD at high temperature. *Physical Review D*, 53:3421–3437, 1996.

[17] D. Kutasov; F. Larsen. Partition sums and entropy bounds in weakly coupled CFT. *Journal of High Energy Physics*, 1, 2001. arXiv:0009.244.

[18] D. Klemm; A. C. Petkou; G. Siopsis. Entropy bounds, monotonicity properties and scaling in CFTs. *Nuclear Physics B*, 601:380–394, 2001. arXiv:0101.076.

[19] R. G. Cai. Cardy-Verlinde formula and AdS black holes. *Physical Review D*, 63, 2001. arXiv:0102.113.

[20] F. L. Lin. Casimir effect of graviton and the entropy bound. *Physical Review D*, 63, 2001. arXiv:0010.127.

[21] R. Brustein; S. Foffa; G. Veneziano. CFT, holography and causal entropy bound. *Physics Letters B*, 507:270–276, 2001. arXiv:0101.083.

[22] V. Ivrii. Second term of the spectral asymptotic expansion of the Laplace-Beltrami operator on manifolds with boundary. *Funcitonal Analysis and its Applications*, 14(2):98–106, 1980.

[23] Y. Netrusov; Y. Safarov. Weyl asymptotic formula for the Laplacian on domains with rough boundaries. *Communications in Mathematical Physics*, 253(2):481–509, 2005.

[24] M. C. Gutzwiller. *Chaos in Classical and Quantum Mechanics*. Springer, 1990.

[25] S. Mukherji; S. S. Pal. Logarithmic corrections to black hole entropy and AdS/CFT correspondence. *Journal of High Energy Physics*, 26, 2002. arXiv:0205.164.

[26] S. Das; P. Majumdar; R. K. Bhaduri. General logarithmic corrections to black hole entropy. *Classical and Quantum Gravity*, 19(9):2355–2368, 2002. arXiv:0111.001.

[27] D. Tong. *Lectures on string theory*. University of Cambridge, 2012.

[28] B. Swingle. Conformal field theory on the Fermi surface. 2010. arXiv:1002.4635.

[29] G. T. Horowitz; H. Ooguri. Spectrum of large N gauge theory from supergravity. *Physical Review Letters*, 80:4116–4118, 1998. arXiv:9802.116.

[30] A. Higuchi. Symmetric tensor spherical harmonics and their application to the de-Sitter group SO(N, 1). *Journal of Mathematical Physics*, 28, 1986.